

**Optimization of kojic acid production from  
*Aspergillus flavus* and *aspargillus oryzae* in  
Solid state fermentation**

---



By:

Afia Zahoor

ID: 15001254006

Supervisor:

Dr. Tanveer Akber

---

**Department of Life Sciences  
School of Science  
University of Management and Technology  
Lahore, Pakistan**

**2017**

**OPTIMIZATION OF KOJIC ACID PRODUCTION FROM  
*ASPERGILLUS FLAVUS* AND *ASPARGILLUS ORYZAE* IN  
SOLID STATE FERMENTATION**

---

Submitted to University of Management and Technology Lahore

In partial fulfillment of the requirements

For the award of degree of

**MS**

**BIOTECHNOLOGY**

By

Afia Zahoor

ID

15001254006

Session: 2015-2017

**Department of Life Sciences  
School of Science  
University of Management and Technology,  
Lahore, Pakistan**



**In the name of**

**ALLAH,**

**The most Compassionate,**

**The most Merciful**

**“Research is what I’m doing when  
I don’t know what I’m doing”**

**- *Wernher von Braun***

## **DECLARATION**

---

I Afia Zahoor D/O Zahoor ul Haq ID: 15001254006 Session 2015-2017 have declare that the matter printed in the thesis dissertation titled “**Optimization of Kojic Acid production from *Aspergillus flavus* and *Aspergillus Oryzae* in solid state fermentation**” which I here submit for the research qualification **M.S Degree In Biotechnology** to the Department of Life Sciences, School of Science, University of Management and Technology, Lahore, Pakistan is apart from the recognized assistance of my supervisors, my own work and has not been previously submitted as research work, thesis or publication in any form in any University, research institution etc. in Pakistan or abroad to obtain a research diploma or degree.

Dated: \_\_\_\_\_

---

(Afia Zahoor)

# DEDICATION

I dedicate this thesis in the honor of our beloved Prophet Hazrat Muhammad (P.B.U.H) and my parents especially to my father (Zahoor ul Haq) without your motivation love and prayers, I could never be able to advance in life.

## ACKNOWLEDGEMENTS

All the praises are for **ALMIGHTY ALLAH**, who is the most beneficent and merciful. I want to thank **ALMIGHTY ALLAH**, who created this universe and bestowed the mankind and me with knowledge and wisdom. HE invigorates me with the ability to accomplish this task of research work and to contribute a drop to the existing ocean of scientific knowledge.

Trembling lips and wet eyes to praise for **PROPHET MUHAMMAD (S.A.W.W)** the city of knowledge, torch of guided who guided His ummah to the knowledge from cradle to grave. The preaching and examples set by our **HOLY PROPHET MUHAMMAD (S.A.W.W)** will remain forever source of guidance for Muslims and whole humanity.

I wish to express my sincere thanks to, **Dr. Tanveer Akber** Assistant Professor, Life Sciences Department University of Management and Technology Lahore, for providing me with a great opportunity to do my research and also for providing me with all facilities during my research work.

It's a truth that I really have no words to say thanks to, **Dr. Yasir Saleem** Senior Scientific Officer at **PCSIR Lahore**, and **Mrs. Shaista Yasir** who always encouraged and helped me during the course of this research work at meat lab PCSIR Lahore, for the continuous encouragement and also for helpful suggestions and deeds during my research work. I will never be able to say thanks for his guidance not only during my research work but also during my study years.

I would also like to say special thanks to **Sir Arshad Sindhu** Leather department PCSIR. I would like to say special thanks to my mother **Nighat Zahoor** and my friend **Muhammad Bilal** and **Samiya Yaqub** for moral support and encouragement.

**AFIA ZAHOOR**

## ABSTRACT

Kojic Acid is a fungal secondary metabolite that can be obtained naturally by the fermentation of fungal species. *Aspergillus* is considered to be the best specie for the production of kojic acid. Kojic Acid has numerous uses in health, cosmetics and pharmaceutical industries. In order to get the highest yield of kojic aid we have used local strains of *Aspergillus* that was *Aspergillus flavus* and *Aspergillus oryzae*. Both strains were giving yield of Kojic Acid in solid state fermentation. Glucose and sucrose were used as carbon source, ammonium nitrate used as a nitrogen source at optimized pH and temperature. Given temperature was 25°C, 30°C, 35°C and 40°C, while given pH was 3.5, 4.5, 5.5, 6.5, 7.5, and 8.5. Highest yield of Kojic Acid was obtained by *Aspergillus flavus* and *Aspergillus oryzae* at 30°C. *Aspergillus flavus* shown best result at pH 4.5 and 3.5 was the optimized pH for *Aspergillus oryzae*. Here we have get the highest yield of Kojic Acid by using local species of *Aspergillus* by optimizing the conditions.

## Contents

Chapter 1 .....	4
INTRODUCTION.....	5
1.1 Kojic Acid .....	5
1.2 Formation of Kojic Acid.....	5
1.3 Chemical properties.....	5
1.4 Physical properties .....	6
1.5 Use of Kojic Acid in food industry.....	6
1.6 Pharmaceutical industry .....	7
1.7 The role of Kojic Acid in cosmetic industry .....	8
1.8 Food industry.....	8
1.9 <i>Aspergillus</i> in the production of kojic acid.....	9
CHAPTER NO: 2.....	10
Review of literature.....	10
Chapter 2 Review of literature.....	10
Literature review .....	10
Chapter 3 Materials and methods.....	20
3.1 STRAINS AND CULTURE.....	31
3.2 MAINTANCE OF THE CULTURE.....	31
3.2.1 Microorganism.....	31
3.3 Optimization of fermentation procedure.....	32
3.3.1 Media .....	32
3.3.2 pH.....	32
3.3.3 Temperature .....	34
Chapter 4 Results and discussion.....	34
4.1 Analysis of Kojic Acid .....	34
4.2 Extraction of kojic acid.....	35
4.3 Thin layer chromatography.....	36
4.4 FTIR.....	37

4.5 Effect of pH on the growth of <i>A.flavus</i> .....	40
4.6 Effect of temperature on the growth of <i>A.flavus</i> .....	42
4.7 <i>Aspergillus oryzae</i> optimization on different pH.....	45
4.8 Effect of temperature on the growth of <i>A.oryzae</i> .....	48
4.9 Discussion.....	51
References.....	54

# List of tables

4.1 Concentrations of Kojic Acid produced by <i>Aspergillus flavus</i> at different pH in static media, on 10th day.....	41
4.2 Concentrations of Kojic Acid produced by <i>Aspergillus flavus</i> at different pH in static media, on 20th day.....	42
4.3 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different temperatures in Static media, on 10th day.....	44
4.4 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different temperatures in static media, on 20th day.....	45
4.5 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 10th day.....	47
4.6 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 20th day.....	48
4.7 Concentrations of Kojic Acid produced by <i>Aspergillus oryzae</i> at different temperatures in static media, on 10th day.....	50
4.8 Concentrations of Kojic Acid produced by <i>Aspergillus oryzae</i> at different temperatures in static media, on 20th day.....	51

## List of figures

3.1 Growth of <i>Aspergillus flavus</i> at pH 4.5-8.5 on solid state media.....	32
3.2 Growth of <i>Aspergillus flavus</i> and <i>Aspergillus oryzae</i> at temperature 25 <sup>o</sup> C-30 <sup>o</sup> C.....	33
4.1 Bentley Method for the confirmation of Kojic Acid.....	35
4.2 Crystals of Kojic Acid in petri dish.....	36
4.3 Confirmation of the presence of Kojic Acid by thin layer chromatography.....	37
4.4 Concentration of original Kojic Acid from FTIR.....	38
4.5 Concentration of Kojic Acid by <i>Aspergillus flavus</i> fungus through FTIR.....	39
4.6 Concentration of Kojic Acid by <i>Aspergillus oryzae</i> fungus through FTIR.....	40
4.7 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different pH.....	42
4.8 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different pH in static media, on 20 <sup>th</sup> day.....	43
4.9 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different temperatures in static media, on 10 <sup>th</sup> day.....	45
4.10 Concentration of Kojic Acid produced by <i>Aspergillus flavus</i> at different temperatures in static media, on 20th day.....	46
4.11 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 10th day.....	48
4.12 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 20th day.....	49
4.13 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 10th day.....	51
4.14 Concentration of Kojic Acid produced by <i>Aspergillus oryzae</i> at different pH in static media, on 20th day.....	52

# **CHAPTER NO: 1**

## **INTRODUCTION**

# Chapter 1

## INTRODUCTION

### 1.1 Kojic acid

Kojic Acid is a secondary fungal metabolite that is obtained from fungus species. These species are *Aspergillus oryzae* and *Aspergillus flavus*. It is an organic acid. Both are considered as best fungal strain to produce the Kojic Acid (Rosfarizan *et al.*, 2010). To produce Kojic Acid biologically active strains of fungus undergo aerobic fermentation by using different substrates (El-Aasar, 2006; El-Kady *et al.*, 2007; Wakisaka *et al.*, 1998). They are producing Kojic Acid as a metabolic by product of fermentation. They are also producing molds. These molds are very useful for the production of fermented products like distilled liquor, sweet beverage, and sweet alcoholic seasoning that is most commonly used in Japan (Niwa *et al.*, 2010).